

[Document Name] Application for Patent

[Reference Number] B00-0196

[Date] May 23, 1997

[Addressed to] Commissioner of Patent Office

[IPC] C12P 41/00

[Title of Invention] ORGANIC ANION TRANSPORTER AND GENE  
CODING FOR THE SAME

[Number of Claims] 17

[Inventor]

[Domicile] 1-23-7, Yoshinodai, Sagamihara-shi,  
Kanagawa

[Name] Hitoshi ENDO

[Inventor]

[Domicile] 214-102, Midori-cho, Hachioji-shi,  
Tokyo

[Name] Yoshikatsu KANAI

[Inventor]

[Domicile] 1-10-47, Sakae-cho, Tachikawa-shi, Tokyo

[Name] Takashi SEKINE

[Inventor]

[Domicile] 3-42-4-301, Shimorenjaku, Mitaka-shi,  
Tokyo

[Name] Makoto HOSOYAMADA

[Applicant]

[Domicile] 1-23-7, Yoshinodai, Sagamihara-shi,

Kanagawa

[Name] Hitoshi ENDO

[Attorney]

[Identification Number] 100076923

[Patent Attorney]

[Name] Shigeo MINOURA

[Phone Number] 06-300-2726

[Indication of Fee]

[Deposit Number] 016322

[Fee] 21000

[List of Document(s)]

[Name of Document]	Specification	1
--------------------	---------------	---

[Name of Document]	Drawings	1
--------------------	----------	---

[Name of Document]	Abstract	1
--------------------	----------	---

[Proof] required

[Designation of Document] Specification

[Title of the Invention] Organic Anion Transporter and Gene Thereof

[Claims]

[Claim 1] Protein selected from the following (A), (B), (C) and (D).

(A) Protein comprising an amino acid sequence represented by SEQ ID No. 1;

(B) Protein comprising an amino acid sequence where one or several amino acid(s) is/are deleted from, substituted for or added to the amino acid sequence represented by SEQ ID No. 1 and having a capability of transporting an organic anion;

(C) Protein comprising an amino acid sequence represented by SEQ ID No. 2; and

(D) Protein comprising an amino acid sequence where one or several amino acid(s) is/are deleted from, substituted for or added to the amino acid sequence represented by SEQ ID No. 2 and having a capability of transporting an organic anion.

[Claim 2] The protein according to claim 1, wherein it is derived from human being.

[Claim 3] The protein according to claim 1, wherein it is derived from rat.

[Claim 4] The protein according to claim 1, wherein it is derived from renal tissues.

[Claim 5] Gene coding for the protein mentioned in

claim 1.

[Claim 6] Gene comprising DNA selected from the following (a), (b), (c) and (d).

(a) DNA comprising a base sequence represented by SEQ ID No. 1;

(b) DNA hybridizing to the DNA comprising the base sequence represented by SEQ ID No. 1 under a stringent condition and coding for protein which has a capability of transport of an organic anion;

(c) DNA comprising a base sequence represented by SEQ ID No. 2;

(d) DNA hybridizing to the DNA comprising the base sequence represented by SEQ ID No. 2 under a stringent condition and coding for protein which has a capability of transport of an organic anion.

[Claim 7] The DNA according to claim 6, wherein it is derived from human being.

[Claim 8] The Gene according to claim 6, wherein it is derived from rat.

[Claim 9] The Gene according to claim 6, wherein it is derived from renal tissues.

[Claim 10] Plasmid containing a region coding for the gene mentioned in any of claims 5 to 9 or protein in the said gene.

[Claim 11] The plasmid according to claim 10, wherein

it is an expression plasmid.

[Claim 12] A host cell which is transformed by the plasmid mentioned in claim 10.

[Claim 13] Nucleotide containing a partial sequence of continuous 14 or more bases in the base sequence represented by SEQ ID No. 1 or No. 2 or a complementary sequence thereof.

[Claim 14] The nucleotide according to claim 13, wherein it is used as a probe for the detection of gene which codes for a protein having a capability of transport of an organic anion.

[Claim 15] The nucleotide according to claim 13, wherein it is used for modulation of expression of gene which codes for a protein having a capability of transport of an organic anion.

[Claim 16] Antibody to the protein mentioned in any of claims 1 to 4.

[Claim 17] A method of using the protein mentioned in any of claims 1 to 4 for the test of action of a test substance as a substrate for the capability of the said protein to transport an organic anion.

[Detailed Description of the Invention]

[0001]

[Technical Field to which the Invention Belongs]

The present invention relates to a gene participating in transport of an organic anion and also to a polypeptide for

which the gene codes.

[0002]

[Prior Art]

Kidney plays an important role for excretion of xenobiotics and drugs outside the body. Anionic drugs are excreted into urine from urinary tubule near the kidney by a route mediated by a carrier. Excretion of an organic anion as such starts by the fact that the urinary tubule cells take up the organic anion from the blood near the urinary tubule via the basolateral membrane.

[0003]

With regard to the uptake of an organic anion by the basolateral membrane, investigations have been carried out by an experiment by means of a perfusion of excised organ or an isolated cell membrane vehicle system using, for example, a p-aminobenzoate as an organic anion of the substrate. During the investigations, it has been believed that an organic anion transporter is participated in the uptake of an organic anion and that the uptake of an organic anion in a basolateral membrane is mediated by an exchange transporter for organic anion with dicarboxylic acid.

[0004]

However, in the conventional means, it is difficult to analyze the details of the transport mechanism in urinary tubules such as a network of transport among transporters and an

interaction among drugs during the process of renal excretion and there has been a demand that gene of an organic anion transporter is isolated so as to make the detailed function analysis possible.

[0005]

With regard to an organic anion transporter gene expressed in liver, various molecular species have been cloned (Hagenbuch, et al., *Proc. Natl. Acad. Sci. USA*, volume 88, page 10629, 1991; Jacquemin, et al., *Proc. Natl. Acad. Sci. USA*, volume 91, page 133, 1994; Shi, et al., *J. Biol. Chem.*, volume 270, page 25591, 1995; and Kanai, et al., *Am. J. Physiol.*, volume 270, page F319, 1996). A genetic cloning of OCT1 which is one of the organic cation transporters expressed in kidney and liver has been also reported (Grundemann, et al., *Nature*, volume 372, page 549, 1994).

[0006]

As to a transporter for dicarboxylic acid, a genetic cloning of sodium-dependent dicarboxylate transporter (NaDC-1) in kidney has been also reported (Pajor, et al., *J. Biol. Chem.*, volume 270, page 5779, 1995).

[0007]

Recently, as an analogous gene of sodium-independent rat liver organic anion transporter (oatp), cloning of gene of an organic anion transporter OAT-K1 localized in renal urinary tubules of rat has been reported (Saito, et al., *J. Biol. Chem.*,

volume 270, page 20719, 1996). However, with regard to the OAT-K1, it has not been confirmed yet that its transport mechanism is due to an exchange transport between an organic anion and a dicarboxylic acid.

[0008]

[Problems that the Invention is to Solve]

An object of the present invention is to provide a novel organic anion transporter gene participating in an organic anion transport in the kidney and also to provide an organic anion transporter which is a polypeptide for which the gene codes. Other objects will be apparent from the following descriptions.

[0009]

[Means for Solving the Problems]

The present inventors have cloned the gene of a novel protein having a capability of transporting an organic anion from renal cells of rats and further cloned a human homologous gene (homolog). They have moreover succeeded in confirming the transporting ability of an organic anion by expressing the product of those genes in oocytes of *Xenopus* whereupon the present invention has been achieved.

[0010]

Thus, the present invention relates to a protein selected from the following (A), (B), (C) and (D).

(A) Protein comprising an amino acid sequence represented by SEQ ID No. 1;



(B) Protein comprising an amino acid sequence where one or several amino acid(s) is/are deleted from, substituted for or added to the amino acid sequence represented by SEQ ID No. 1 and having a capability of transporting an organic anion;

(C) Protein comprising an amino acid sequence represented by SEQ ID No. 2; and

(D) Protein comprising an amino acid sequence where one or several amino acid(s) is/are deleted from, substituted for or added to the amino acid sequence represented by SEQ ID No. 2 and having a capability of transporting an organic anion.

[0011]

The present invention further relates to a gene comprising DNA selected from the following (a), (b), (c) and (d).

(a) DNA comprising a base sequence represented by SEQ ID No. 1;

(b) DNA hybridizing to the DNA comprising the base sequence represented by SEQ ID No. 1 under a stringent condition and coding for protein which has a capability of transport of an organic anion;

(c) DNA comprising a base sequence represented by SEQ ID No. 2;

(d) DNA hybridizing to the DNA comprising the base sequence represented by SEQ ID No. 2 under a stringent condition and coding for protein which has a capability of transport of an organic anion.

[0012]

In living bodies, the novel protein having a capability of transporting an organic anion according to the present invention or, in other words, an organic anion transporter (OAT1) is mostly expressed in urinary tubules of the kidney.

[0013]

Further, an organic anion transporting ability (uptake of an organic anion into expressed cells) of the organic anion transporter OAT1 is activated by the presence of a dicarboxylic acid. It is therefore likely to be a transporter which carries out an exchange transport between an organic anion and a dicarboxylic acid. Further, in the exchange transport, it is likely that the dicarboxylic acid which comes outside the cells in exchange for the organic anion by OAT1 is taken up into the cells by a sodium-dependent dicarboxylic acid transporter (NaDC-1) for recycling.

[0014]

Furthermore, the organic anion transporter OAT1 of the present invention has a substrate selectivity of a very broad range having a capability of transporting (taking up) various drugs having different structures such as cyclic bases, prostaglandins, uric acid, antibiotic substances, nonsteroidal anti-inflammatory agents, diuretics and anti-tumor agents.

[0015]

Still further, it is likely that the organic anion

transporter OAT1 of the present invention has no homology to the already-reported organic anion transporter OAT-K1 from kidney of rats but is an entirely different molecular species.

[0016]

[Mode for Carrying Out the Invention]

SEQ ID No. 1 in the Sequence Listing which will be mentioned later shows an amino acid sequence (551 amino acids) of protein encoded to the full-length cDNA base sequence (about 2.2 kbp) of gene of an organic anion transporter (rat OAT1) derived from kidney of rats and a translation region thereof.

[0017]

SEQ ID No. 2 shows an amino acid sequence (563 amino acids) of protein encoded to the full-length cDNA base sequence (about 2.2 kbp) of gene of an organic anion transporter (human OAT1) derived from kidney of human being and a translation region thereof.

[0018]

With regard to the base sequences or the amino acid sequences shown in the above SEQ ID Nos. 1 and 2, a homology search was carried out for all sequences included in the already-known DNA databases (GenBank and EMBL) and protein databases (NBRF and SWISS-PROT) and, as a result, there was no identical one therein whereupon those sequences are thought to be novel ones.

[0019]

With regard to the protein of the present invention, in addition to that which has an amino acid sequence represented by SEQ ID No.1 or 2, those in which one or more amino acid(s) in the amino acid sequence represented by SEQ ID No. 1 or 2 is/are deleted, substituted or added may be exemplified. Deletion, substitution or addition of amino acid(s) may be within such an extent that an organic anion transport activity is not lost and the number(s) is/are usually from 1 to about 110 or, preferably, from 1 to about 55. Such an amino acid has a homology in terms of an amino sequence to the amino acid sequence represented by SEQ ID No. 1 or 2 to an extent of usually 1-80% or, preferably, 1-90%.

[0020]

With regard to the gene of the present invention, there are exemplified that which contains DNA being able to hybridize to DNA having a base sequence represented by SEQ ID No. 1 or 2 under a stringent condition in addition to that containing DNA having a base sequence represented by SEQ ID No. 1 or 2. The DNA which is able to hybridize as such may be in such an extent that protein for which the DNA is coded has a capability of transporting the organic anion. Such a DNA has a homology of usually 70% or more or, preferably, 80% or more homology to the base sequence represented by SEQ ID No. 1 or 2. Such a DNA includes mutant gene which is found in nature, mutant gene which is artificially modified and homologous gene derived

from other living body.

[0021]

In the present invention, hybridization under a stringent condition is usually carried out in such a manner that hybridization is conducted in a hybridization solution of 5 × SSC or similar salt concentration thereto at the temperature condition of 37 to 42°C for about 12 hours, a preliminary washing is conducted if necessary with a solution of 5 × SSC or similar salt concentration thereto and then a washing is conducted with a solution of 1 × SSC or similar concentration thereto. In order to achieve a higher stringency, it can be done where a washing is conducted in a solution of 0.1 × SSC or similar salt concentration thereto.

[0022]

It is also possible that the organic anion transporter gene according to the present invention is isolated and obtained by means of a screening where tissues or cells of appropriate mammals are used as a gene source. With regard to the mammals, human being may be exemplified in addition to non-human ones such as dogs, cattle, horses, goats, sheep, monkeys, pigs, rabbits, rats and mice.

[0023]

Screening and isolation of the gene may be advantageously carried out by means of an expression cloning, etc.

[0024]

For example, renal tissues of rats are used as a gene source and mRNA (poly(A)<sup>+</sup>RNA) is prepared therefrom. This is fractionated and each of the fractions is introduced into oocytes of *Xenopus* together with cRNA of rat sodium-dependent dicarboxylate transporter (NaDC-1).

[0025]

cDNA of NaDC-1 gene has been reported already (Pajor, et al., *J. Biol. Chem.*, volume 270, page 5779, 1995) and, therefore, it is possible to easily prepare the cDNA of NaDC-1 gene from the sequence information using a PCR or the like. Starting from the resulting NaDC-1 cDNA, it is possible to synthesize RNA (cRNA) (being capped) which is complementary thereto using, for example, T3 or T7 RNA polymerase.

[0026]

With regard to the oocytes into which mRNA and NaDC-1 cRNA are introduced, transport (uptake) of a substrate into the cells is measured using, for example, p-aminohippuric acid (PAH) as a substrate (organic anion) and the fraction of mRNA showing a high uptake is selected whereupon mRNA in OAT1 can be concentrated. A cDNA library is prepared on the basis of the mRNA concentrated as such. cRNA (capped one) is prepared from cDNA of the library, each clone is introduced into oocytes together with NaDC-1 cRNA as same as before and positive clone is selected using the uptake activity of the substrate as an index whereupon it is possible to obtain clone containing cDNA

of the OAT1 gene.

[0027]

As to the resulting cDNA, its base sequence is determined by a conventional method and a translated region is analyzed whereby the protein coded therefor or the amino acid sequence of OAT1 can be determined.

[0028]

The fact that the resulting cDNA is a cDNA of an organic anion transporter gene or, in other words, the gene product coded for cDNA is an organic anion transporter can, for example, be inspected as follows. Thus, cRNA prepared from the resulting OAT1 cDNA is introduced into oocytes to express and an ability of transporting (taking up) the organic anion into the cells can be confirmed as above by measuring the uptake of the substrate into the cells by means of a conventional uptake experiment (Kanai and Hediger, *Nature*, volume 360, pages 467-471, 1992) using an appropriate organic anion as a substrate.

[0029]

It is also possible that the same uptake experiment is applied for the expression cells to investigate the characteristics of OAT1 such as a characteristic that OAT1 conducts an exchange transport with dicarboxylic acid and a substrate specificity of OAT1.

[0030]

When an appropriate cDNA library or genomic DNA library

prepared from a different genetic source is screened using cDNA of the resulting OAT1 gene, it is possible to isolate homologous gene, chromosomal gene, etc. from different tissues and different living bodies.

[0031]

It is further possible to isolate the gene from a cDNA library or a genomic DNA library by a conventional PCR (Polymerase Chain Reaction) using a synthetic primer designed on the basis of the information of the disclosed base sequence of the gene of the present invention (base sequence as shown in SEQ ID No. 1 or 2 or a partial sequence thereof).

[0032]

The DNA library such as a cDNA library and a genomic DNA library can be prepared by a method mentioned, for example, in "Molecular Cloning" (by Sambrook, J., Fritsch, E. F. and Maniatis, T.; published by Cold Spring Harbor Laboratory Press in 1989). Alternatively, when a commercially available library is available, that may be used as well.

[0033]

The organic anion transporter (OAT1) of the present invention may be produced, for example, by means of a gene recombination technique using cDNA coding for the organic anion transporter. For example, DNA (cDNA or the like) coding for the organic anion transporter is incorporated into an appropriate expression vector and the resulting recombinant



DNA can be introduced into an appropriate host cell. Examples of the expression system (a host-vector system) for the production of a polypeptide are expression systems of bacteria, yeasts, insect cells and mammalian cells. In order to obtain a functional protein among them, it is desirable to use insect cells and mammalian cells.

[0034]

For example, in order to express a polypeptide in mammals, DNA coding for an organic anion transporter is inserted into a downstream to an appropriate promoter (such as SV40, LTR promoter and elongation 1 $\alpha$  promoter) in an appropriate expression vector (such as vector of a retrovirus type, papilloma virus vector, vaccinia virus vector and vector of an SV40 type) whereupon an expression vector is constructed. Then appropriate cells are subjected to a transformation using the resulting expression vector and the transformant is incubated in an appropriate medium to give a desired polypeptide. Examples of the mammal cells used as a host cell are cell strains including simian COS-7 cells, Chinese hamster CHO cells, human HeLa cells, primary culture cells derived from kidney tissues, LLC-PK1 cells derived from kidney of pig, OK cells derived from opossum kidney, etc.

[0035]

With regard to DNA which codes for an organic anion transporter OAT1, the cDNA having a base sequence shown in SEQ

ID No. 1 or 2 may be used for example and, in addition, it is not limited to the above-mentioned cDNA but DNA corresponding to the amino acid sequence is designed and the DNA coding for the polypeptide may be used. In that case, 1 to 6 kind(s) of codon has/have been known for coding for each amino acid and, although any codon may be selected for use, a sequence having higher expression can be designed when, for example, frequency of use of codon by the host utilized in the expression is taken into consideration. DNA having a designed base sequence can be obtained by chemical synthesis of DNA, fragmentation and bonding of the above-mentioned cDNA, a partial modification of the base sequence, etc. Artificial partial modification of and introduction of variation into a base sequence can be carried out by a site-specific mutagenesis (Mark, D. F., et al., *Proceedings of National Academy of Sciences*, volume 81, pages 5662 to 5666 (1984)), etc. utilizing a primer comprising a synthetic oligonucleotide coding for the desired modification.

[0036]

The nucleotide (oligonucleotide or polynucleotide) which hybridizes to the organic anion transporter gene of the present invention under a stringent condition can be used as a probe for detecting the organic anion transporter gene and, moreover, it can be used as antisense oligonucleotide, ribozyme, decoy, etc. for modulation of expression of the organic anion

transporter. With regard to such a nucleotide, a nucleotide containing a partial sequence of usually not less than continuous 14 bases in the base sequence represented by SEQ ID No. 1 or 2 or a complementary sequence thereof may be used for example and, in order to hybridize more specifically, longer sequence such as not less than 20 bases or not less than 30 bases may be used as a partial sequence.

[0037]

It is also possible that the organic anion transporter of the present invention or a polypeptide having the immunological homology thereto is used to obtain an antibody thereto and the antibody is able to be utilized for detection, purification, etc. of the organic anion transporter. The antibody can be manufactured using the organic anion transporter of the present invention or a fragment thereof or a synthetic peptide having a partial sequence thereof as an antigen. Polyclonal antibody can be manufactured by a conventional method in which an antigen is inoculated to a host animal (such as rat and rabbit) and immune serum is recovered while monoclonal antibody can be manufactured by a conventional way such as a hybridoma method.

[0038]

The present invention will now be further illustrated by way of the following Examples although those Examples do not limit the present invention.

[0039]

In the following Examples, each operation is carried out, unless otherwise mentioned, by a method mentioned in "Molecular Cloning" (by Sambrook, J., Fritsch, E. F. and Maniatis, T.; published by Cold Spring Harbor Laboratory Press in 1989) or according to the direction for use of the commercially available product when a commercially available reagent or kit is used.

[0040]

[Examples]

Example 1. Cloning of rat organic anion transporter

(1) Isolation of rat dicarboxylate transporter cDNA and preparation of cRNA

cDNA library was prepared from rat poly(A)<sup>+</sup> RNA using a kit for the synthesis of cDNA (trade name: SuperScript Choice System; manufactured by Gibco) and was integrated into a site of phage vector  $\lambda$  Ziplox (manufactured by Gibco) cleaved by a restriction enzyme EcoRI. Segment corresponding to the bases (from the 1323rd to the 1763rd) of rabbit sodium-dependent dicarboxylic acid transporter NaDC-1 gene (Pajor, et al., *J. Biol. Chem.*, volume 270, page 5779, 1995) was labeled with <sup>32</sup>P-dCTP and, using it as a probe, a cDNA library of rat was screened by PCR method. Hybridization was carried out for one night in a solution for hybridization of 37°C and the filter membrane was washed at 37°C with 0.1 × SSC/0.1% SDS. As to the solution for hybridization, a buffer of pH 6.5 containing 5

x SSC, 3x Denhard's solution, 0.2% of SDS, 10% of dextran sulfate, 50% of formamide, 0.01% of Antiform B (trade name; manufactured by Sigma) (a defoaming agent), 0.2 mg/ml of DNA modified by salmon sperm, 2.5 mM of sodium pyrophosphate and 25 mM of MES was used. The cDNA moiety integrated in  $\lambda$  Ziplox phage was integrated into plasmid pZL1 for the determination of a base sequence and further subcloned to a plasmid pBluescript IISK (manufactured by Stratagene).

[0041]

cRNA (RNA complementary to cDNA) was prepared from a plasmid containing cDNA of the above-prepared rat dicarboxylate transporter using a T7 RNA polymerase.

[0042]

The resulting cRNA was injected into oocytes of *Xenopus* according to a method by Kanai, et al. (Kanai and Hediger, *Nature*, volume 360, pages 467 to 471, 1992) and, using the oocytes, an uptake experiment was carried out using glutaric acid as a substrate. In the experiment, radioactively labeled substrate ( $^{14}\text{C}$ -glutaric acid) was used. As a result, uptake of glutaric acid in a sodium-dependent manner was noted and it was confirmed that the cloned cDNA was that of rat dicarboxylate transporter gene (Fig. 1).

[0043]

(2) Cloning of rat kidney organic anion transporter OAT1  
It was carried out by an expression cloning method as

follows according to a method by Kanai, et al. (Kanai and Hediger, *Nature*, volume 360, pages 467 to 471, 1992).

[0044]

Rat kidney poly (A)<sup>+</sup> RNA (400 µg) was fractionated by a gel electrophoresis.

[0045]

Each fraction obtained by the fractionation was injected into oocytes together with cRNA of the rat dicarboxylate transporter obtained in the above (1). The oocytes used were those which were previously incubated for 2 hours in a sodium uptake solution containing 1 mM of glutaric acid as a substrate (96 mM sodium chloride, 2 mM potassium chloride, 1.8 mM calcium chloride, 1 mM magnesium chloride and 5 mM HEPES; pH 7.4).

[0046]

The oocytes into which RNA was injected were subjected to an experiment for the uptake of substrate using p-aminohippurate (hereinafter, abbreviated as PAH) as a substrate according to a method by Kanai, et al. (Kanai and Hediger, *Nature*, volume 360, pages 467 to 471, 1992). Oocytes were incubated for 1 hour in a sodium uptake solution containing <sup>14</sup>C-PAH (50 µM) as a substrate and containing no glutaric acid and the uptake rate of the substrate was measured by way of counting the radioactivity taken up into the cells. It was incidentally confirmed that, in this system, uptake of PAH was not noted in oocytes where only poly(A)<sup>+</sup> RNA (mRNA) of kidney

of rat was injected and in oocytes where only cRNA of rat dicarboxylate transporter was injected while, in oocytes where both poly(A)<sup>+</sup> RNA of kidney of rat and dicarboxylate transporter cRNA of rat were injected, uptake of PAH was noted (Fig. 2).

[0047]

Among the RNA fractions, there was selected a fraction where the oocytes into which RNA was injected showed the highest uptake rate of PAH. With regard to poly(A)<sup>+</sup> RNA (1.8 to 2.4 kb) of this fraction, a cDNA library was prepared using a kit for cDNA synthesis and plasmid cloning (trade name: Superscript Plasmid Systems; manufactured by Gibco). Those DNAs were integrated with the sites recognizing SalI and NotI which were restriction enzymes of plasmid pSPORT1 (manufactured by Gibco) and the resulting recombinant plasmid DNA was introduced into competent cells (trade name: Electro Max DH10B Competent Cell; manufactured by Gibco BRL) of DH10B strain of *Escherichia coli*. The resulting transformant was incubated on a nitrocellulose membrane to give about 500 colonies per plate. Plasmid DNAs were prepared from those colonies and cleaved with a restriction enzyme NotI. Capped cRNA was synthesized using the resulting DNAs by means of an *in vitro* transcription.

[0048]

The resulting cRNA (about 10 ng) was injected into oocytes together with cRNA (2 ng) of the rat dicarboxylate transporter obtained in the above (1). Those oocytes were subjected, as

same as above, to a screening for positive clone by carrying out an uptake experiment for PAH. In the screening, investigation was conducted for a group where DNA extracted from a plurality of clones was pooled and, when uptake of p-aminohippuric acid was confirmed in a group, it was further divided into plural groups followed by conducting a screening.

[0049]

As a result of the screening, one positive clone (clone where uptake of a substrate was noted in oocytes into which cRNA was injected) was isolated from 8,000 clones.

[0050]

With regard to the resulting clone or, in other words, the clone containing cDNA of rat dicarboxylate transporter OAT1, its base sequence was determined by a dideoxy method using a kit for the preparation of deleted clone for determining the base sequence (trade name: Kilo-Sequence Deletion Kit; manufactures by Takara Shuzo), synthetic primer and a kit for the determination of base sequence (trade name: Sequenase ver. 2.0; manufactured by Amersham).

[0051]

As a result, there was obtained a base sequence of cDNA of the rat dicarboxylate transporter OAT1 gene. Further, the base sequence of cDNA was analyzed by a conventional means whereupon the translation region on cDNA and an amino acid sequence of OAT1 coded thereon were determined.



[0052]

Those sequences were shown in SEQ ID No. 1 of the Sequence Listing which will be given later.

[0053]

As a result of analysis of an amino acid sequence of OAT1 by a Kyte-Doolittle hydropathy analysis (hydrophobic plot), 12 membrane-spanning domains were predicted as shown in Fig. 3. In addition, 5 sugar chain-added sites were predicted in the first hydrophilic loop. In a loop of a hydrophilic group of the 6th and 7th transmembrane domains, there were 4 sites which were thought to be proteinkinase C-dependent phosphorylated sites.

[0054]

(3) Expression of OAT1 gene in various tissues (analysis by a northern blotting)

A full-length cDNA of rat OAT1 gene was labeled with  $^{32}\text{P}$ -dCTP and, using it as a probe, a northern blotting was carried out as follows to RNA extracted from various tissues of rat. Poly(A)<sup>+</sup> RNA (3  $\mu\text{g}$ ) was subjected to electrophoresis using 1% agarose/formaldehyde gel and transferred to a nitrocellulose filter. This filter was subjected to hybridization at 42°C for one night using a hybridization solution containing a full-length OAT1 cDNA labeled with  $^{32}\text{P}$ -dCTP. The filter was washed at 65°C with 0.1  $\times$  SSC containing 0.1% of SDS.

[0055]

As a result of the northern blotting (Fig. 4), in kidney, two bands corresponding to about 2.9 kb and 3.9 kb/4.2 kb were detected whereby expression was noted. In cortex of kidney and outer layer of medulla, expressed amount of OAT1 mRNA was large while it was small in inner layer of medulla.

[0056]

When exposed to light for longer time, a slight band was detected at about 2.4 kb in brain while, in all other tissues, no band was detected and no expression was noted.

[0057]

(4) Expression of OAT1 gene in renal tissues (analysis by an *in situ* hybridization)

An *in situ* hybridization was carried out as follows. Thus, after kidney of rat was fixed by perfusion with 4% paraformaldehyde, it was finely cut and further fixed with 4% paraformaldehyde. The resulting kidney of rat was sliced in a thickness of 5  $\mu$ m and the resulting slices were used for an *in situ* hybridization.

[0058]

From a full-length OAT1 cDNA were synthesized <sup>35</sup>S-labeled sense cRNA and antisense cRNA using T7 or T3 RNA polymerase and they were used as probes. The slice was subjected to hybridization with the probe for one night using a hybridization solution followed by washing with 0.1  $\times$  SSC at 37°C for 30 minutes.

[0059]

It was shown as a result of the *in situ* hybridization that, in the layered sites of kidney of rat, OAT1 mRNA was expressed in cortex of kidney and outer layer of medulla, particularly at the part of medullar ray of cortex. In the inner layer of medulla, no expression was detected. The result shows that the organic anion transporter OAT1 was most abundantly expressed in the middle part of proximal convoluted tubule.

[0060]

Example 2. Characterization of organic anion transporter OAT1

(1) Influence of glutaric acid on transport activity of OAT1

In an uptake experiment of PAH by oocytes into which rat OAT1 gene cRNA was injected, influence of preincubation with glutaric acid was investigated.

[0061]

The uptake experiment of PAH was carried out as follows according to a method described in (2) of the above Example 1. Thus, oocytes into which rat OAT1 gene cRNA or rat OAT1 gene cRNA and rat NaDC-1 cRNA was/were injected were subjected to a preincubation for 2 hours in a sodium uptake solution to which 1 mM glutaric acid was or was not added, then  $^{14}\text{C}$ -PAH was added, the mixture was incubated at room temperature for 1 hour and uptake of the substrate labeled with radioactivity was measured.

[0062]

As a result, uptake of PAH increased when the oocytes were subjected to a pretreatment with 1 mM glutaric acid (Fig. 5). When oocytes where rat dicarboxylate transporter and OAT1 were expressed were subjected to a pretreatment with glutaric acid, further increase in uptake of  $^{14}\text{C}$ -p-aminohippurate was noted. The effect of glutaric acid noted in such a result shows the dependency of PAH uptake on the dicarboxylate concentration in cells whereby it is likely that OAT1 is an exchange transporter for organic anion and dicarboxylic acid.

[0063]

(2) Dependency of transport activity of OAT1 on salt

In an uptake experiment of PAH by oocytes into which rat OAT1 gene cRNA was injected, influence of a salt which was added to a medium was investigated.

[0064]

An uptake experiment for PAH was carried out according to a method mentioned in the above (1) using oocytes into which rat OAT1 gene cRNA was injected. With regard to an uptake solution, however, a choline chloride uptake solution (a sodium solution of 96 mM sodium chloride was changed to 96 mM choline chloride followed by adjusting to pH 7.4) when influence of addition of choline chloride ion was added was checked instead of sodium uptake solution.

[0065]

The result (Fig. 6) was that, even when sodium outside the cells was substituted with choline, no influence was noted on the uptake of PAH. Therefore, it was noted that OAT1 was a transporter which acted in an independent basis to sodium ion.

[0066]

(3) Michaelis-Menten's dynamic test for OAT1

A Michaelis-Menten's dynamic test was carried out for an organic anion transporter. Changes in the uptake rate of PAH by the changes in concentration of the substrate PAH were checked whereupon a Michaelis-Menten's dynamic test for organic anion transporter was carried out.

[0067]

An uptake experiment for PAH was carried out according to the method mentioned in (1) using oocytes into which rat OAT1 gene cRNA was injected. Uptake of  $^{14}\text{C}$ -PAH was measured for 3 minutes however. The result (Fig. 7) was that  $K_m$  value was about  $14.3 \pm 2.9 \mu\text{M}$ .

[0068]

The  $K_m$  value was similar to the  $K_m$  value ( $80 \mu\text{M}$ ) for an organic anion transport system at the side of base reported in an *in vivo* system already (Ulrich, et al., *Am. J. Physiol.*, volume 254, pages F453 to 462, 1988).

[0069]

(4) Substrate selectivity of OAT1 (Inhibiting test by

addition of drug)

In a PAH uptake experiment using oocytes into which rat OAT1 gene cRNA was injected, influence of addition of various drugs to the system was checked.

[0070]

The PAH uptake experiment was carried out according to the method mentioned in the above (1) using oocytes into which rat OAT1 gene cRNA was injected. However, a sodium uptake solution was used and uptake of PAH was measured in the presence and absence of various compounds (non-labeled) in an amount of 2 mM.

[0071]

The result (Fig. 8) was that a cis-inhibiting effect was observed by addition of a drug which was unrelated to structure. Cefaloridin (antibiotic substance of a  $\beta$ -lactam type), nalidixic acid (old quinolone), furosemide and ethacrynic acid (diuretics), indomethacin (nonsteroidal anti-inflammatory agent), probenecid (agent for excretion of uric acid) and valproic acid (anti-epilepsy agent) strongly inhibited the uptake of  $^{14}\text{C}$ -p-aminohippurate mediated by OAT-1 (85%>). Methotrexate which is an anti-tumor agent inhibited the uptake of PAH in a medium degree. Endogenous compounds such as prostaglandin E2, c-AMP, c-GMP and uric acid also inhibited the uptake of PAH.

[0072]

(5) Substrate-specificity of OAT1 (Uptake test where various kinds of anionic substances were used as substrates)

Uptake by OAT1 was investigated using various kinds of anionic substances as substrates.

[0073]

The PAH uptake experiment was carried out according to the method mentioned in the above (1) using oocytes into which rat OAT1 gene cRNA was injected. As to the substrate however, various compounds labeled with radioactivity were used in place of  $^{14}\text{C}$ -PAH.

[0074]

The result (Fig. 9) was that uptake to oocytes was noted when methoxalate ( $^3\text{H}$ -labeled substance), c-AMP ( $^3\text{H}$ -labeled substance), c-GMP ( $^3\text{H}$ -labeled substance), prostaglandin ( $^3\text{H}$ -labeled substance), uric acid ( $^{14}\text{C}$ -labeled substance) and  $\alpha$ -ketoglutaric acid ( $^{14}\text{C}$ -labeled substance) were used as substrates. On the other hand, no uptake was noted in the case of TEA ( $^{14}\text{C}$ -labeled substance) and taurocholic acid.

[0075]

Example 3. Cloning of human organic anion transporter

The cDNA fragment of rat OAT1 gene obtained in (2) of Example 1 was labeled and was used as a probe for the screening of human cDNA library. With regard to the human cDNA library, there was used a human cDNA library prepared by using human kidney poly(A)<sup>+</sup> RNA (manufactured by Clontec) as a gene source.

[0076]

With regard to the resulting positive clone, or a clone containing human organic anion transporter (human OAT1) cDNA, its base sequence was determined by the same manner as in Example 1 and the base sequence of the resulting cDNA was analyzed by a conventional method to determine the translation region on cDNA and the amino acid sequence of human OAT1 coded thereon.

[0077]

Sequence of the human OAT1 is shown in SEQ ID No. 2 in the Sequence Listing which will be mentioned later.

[0078]

Homology of rat OAT1 to human OAT1 was about 85% at an amino acid level. Homology at a cDNA level was about 79%.

[0079]

[Advantage of the Invention]

The organic anion transporter OAT1 of the present invention and gene thereof is expected to be useful for clarification of the fate of a drug and the fate of a toxin at a molecular level such as an *in vitro* analysis of excretion of a drug and interaction between drugs. Further, many drugs which are causes for renal failure such as antibiotic substance of  $\beta$  lactam type, diuretics and nonsteroidal anti-inflammatory agents are transported by OAT1 and the cause of induction of renal toxicity by drugs is suggested to be possibly by accumulation due to OAT1 whereby a method for screening a drug



for prevention of renal toxicity using OAT1 is expected to be developed.

[0080]

[SEQUENCE LISTING]

Sequence ID No: 1

Length: 2294

Type: Nucleic acid

Number of Chain: Doubled-Stranded

Topology: Linear

Kind: cDNA to mRNA

Origin

Organism: Rat

Sequence

```
GCTCCAGCAG ACCCTGAAAG CTGAGCTGTC CAGACCCCCG AAGTGAAGAA AAGAGGCGAG 60
GGCAAGGGAG GGCCAGAACC GAGGGAGAGA GAAAGGAGGG GCAGCCCACC AGCCCGCTGT 120
CCTGCCACAG AACCGGCTCA GCTCCAGCTC CAGGAGTCAC TCAGCTGCAG AGGCAGTGGC 180
AGCCCCACTC CTCAGGCAAA GGGCAGCAGA CAGACAGACA GAGGTCCTAG GACTGGAGGT 240
CCTCAGTCAT TGACCACTCA GCCTGGCCCCA GCCCC 275
ATG GCC TTC AAT GAC CTC CTG AAA CAG GTG GGG GGC GTC GGA CGC 320
Met Ala Phe Asn Asp Leu Leu Lys Gln Val Gly Gly Val Gly Arg
1 5 10 15
TTC CAG TTG ATC CAG GTC ACC ATG GTG GTT GCT CCC CTA CTG CTG 365
Phe Gln Leu Ile Gln Val Thr Met Val Val Ala Pro Leu Leu Leu
20 25 30
ATG GCT TCC CAC AAC ACC TTG CAG AAC TTC ACT GCC GCT ATC CCC 410
```

Met	Ala	Ser	His	Asn	Thr	Leu	Gln	Asn	Phe	Thr	Ala	Ala	Ile	Pro		
				35					40					45		
CCT	CAT	CAC	TGC	CGC	CCA	CCT	GCC	AAT	GCC	AAT	CTC	AGC	AAA	GAT	455	
Pro	His	His	Cys	Arg	Pro	Pro	Ala	Asn	Ala	Asn	Leu	Ser	Lys	Asp		
				50					55					60		
GGA	GGT	CTG	GAG	GCC	TGG	CTG	CCC	CTG	GAC	AAG	CAA	GGA	CAA	CCC	500	
Gly	Gly	Leu	Glu	Ala	Trp	Leu	Pro	Leu	Asp	Lys	Gln	Gly	Gln	Pro		
				65					70					75		
GAA	TCG	TGC	CTC	CGC	TTT	ACT	TCC	CCC	CAG	TGG	GGA	CCA	CCC	TTT	545	
Glu	Ser	Cys	Leu	Arg	Phe	Thr	Ser	Pro	Gln	Trp	Gly	Pro	Pro	Phe		
				80					85					90		
TAC	AAT	GGC	ACA	GAA	GCC	AAT	GGC	ACC	AGA	GTC	ACA	GAG	CCC	TGC	590	
Tyr	Asn	Gly	Thr	Glu	Ala	Asn	Gly	Thr	Arg	Val	Thr	Glu	Pro	Cys		
				95					100					105		
ATT	GAT	GGC	TGG	GTC	TAT	GAC	AAC	AGC	ACC	TTC	CCT	TCA	ACC	ATC	635	
Ile	Asp	Gly	Trp	Val	Tyr	Asp	Asn	Ser	Thr	Phe	Pro	Ser	Thr	Ile		
				110					115					120		
GTG	ACT	GAG	TGG	AAC	CTT	GTG	TGC	TCT	CAT	CGG	GCT	TTC	CGC	CAG	680	
Val	Thr	Glu	Trp	Asn	Leu	Val	Cys	Ser	His	Arg	Ala	Phe	Arg	Gln		
				125					130					135		
CTG	GCC	CAG	TCC	CTG	TAC	ATG	GTG	GGA	GTG	CTG	CTG	GGA	GCC	ATG	725	
Leu	Ala	Gln	Ser	Leu	Tyr	Met	Val	Gly	Val	Leu	Leu	Gly	Ala	Met		
				140					145					150		
GTG	TTT	GGC	TAC	CTG	GCG	GAC	AGG	CTG	GGC	CGC	CGG	AAG	GTG	CTG	770	
Val	Phe	Gly	Tyr	Leu	Ala	Asp	Arg	Leu	Gly	Arg	Arg	Lys	Val	Leu		

155	160	165	
ATC TTG AAC TAC CTG CAG ACA GCT GTG TCG GGA ACC TGT GCA GCC			815
Ile Leu Asn Tyr Leu Gln Thr Ala Val Ser Gly Thr Cys Ala Ala			
170	175	180	
TAT GCA CCC AAC TAT ACT GTC TAC TGC GTT TTC CGG CTC CTC TCG			860
Tyr Ala Pro Asn Tyr Thr Val Tyr Cys Val Phe Arg Leu Leu Ser			
185	190	195	
GGC ATG TCT TTG GCT AGC ATT GCA ATC AAC TGC ATG ACA CTA AAT			905
Gly Met Ser Leu Ala Ser Ile Ala Ile Asn Cys Met Thr Leu Asn			
200	205	210	
GTG GAA TGG ATG CCT ATC CAC ACC CGT GCC TAT GTG GGC ACC TTG			950
Val Glu Trp Met Pro Ile His Thr Arg Ala Tyr Val Gly Thr Leu			
215	220	225	
ATT GGC TAT GTC TAC AGC CTG GGC CAG TTC CTC CTG GCT GGC ATC			995
Ile Gly Tyr Val Tyr Ser Leu Gly Gln Phe Leu Leu Ala Gly Ile			
230	235	240	
GCC TAT GCT GTG CCC CAC TGG CGC CAC CTG CAG CTT GTG GTC TCT			1040
Ala Tyr Ala Val Pro His Trp Arg His Leu Gln Leu Val Val Ser			
245	250	255	
GTG CCT TTT TTC ATT GCC TTC ATC TAC TCT TGG TTC TTC ATT GAG			1085
Val Pro Phe Phe Ile Ala Phe Ile Tyr Ser Trp Phe Phe Ile Glu			
260	265	270	
TCA GCC CGC TGG TAC TCC TCC TCA GGA AGG CTG GAC CTC ACC CTC			1130
Ser Ala Arg Trp Tyr Ser Ser Ser Gly Arg Leu Asp Leu Thr Leu			
275	280	285	

CGA GCC CTG CAG AGA GTG GCC CGG ATC AAT GGG AAA CAA GAA GAA	1175
Arg Ala Leu Gln Arg Val Ala Arg Ile Asn Gly Lys Gln Glu Glu	
290 295 300	
GGG GCT AAG CTA AGT ATA GAG GTG CTC CGG ACC AGC CTG CAG AAG	1220
Gly Ala Lys Leu Ser Ile Glu Val Leu Arg Thr Ser Leu Gln Lys	
305 310 315	
GAA CTG ACT CTA AGC AAA GGC CAA GCC TCA GCC ATG GAG CTG CTG	1265
Glu Leu Thr Leu Ser Lys Gly Gln Ala Ser Ala Met Glu Leu Leu	
320 325 330	
CGC TGC CCC ACC CTT CGA CAC CTC TTC CTC TGT CTC TCC ATG CTG	1310
Arg Cys Pro Thr Leu Arg His Leu Phe Leu Cys Leu Ser Met Leu	
335 340 345	
TGG TTT GCC ACT AGC TTT GCC TAC TAC GGG CTG GTC ATG GAC CTG	1355
Trp Phe Ala Thr Ser Phe Ala Tyr Tyr Gly Leu Val Met Asp Leu	
350 355 360	
CAG GGC TTT GGG GTC AGC ATG TAC CTT ATC CAG GTG ATT TTC GGT	1400
Gln Gly Phe Gly Val Ser Met Tyr Leu Ile Gln Val Ile Phe Gly	
365 370 375	
GCC GTG GAC CTG CCT GCC AAG TTT GTA TGC TTC CTA GTC ATC AAC	1445
Ala Val Asp Leu Pro Ala Lys Phe Val Cys Phe Leu Val Ile Asn	
380 385 390	
TCC ATG GGG CGC CGG CCT GCA CAG ATG GCC TCC CTG CTG CTG GCA	1490
Ser Met Gly Arg Arg Pro Ala Gln Met Ala Ser Leu Leu Leu Ala	
395 400 405	
GGC ATC TGC ATC CTG GTG AAT GGC ATA ATA CCG AAG AGC CAT ACG	1535

Gly Ile Cys Ile Leu Val Asn Gly Ile Ile Pro Lys Ser His Thr	
410 415 420	
ATC ATT CGC ACC TCC CTG GCT GTG CTA GGG AAG GGC TGC CTG GCT	1580
Ile Ile Arg Thr Ser Leu Ala Val Leu Gly Lys Gly Cys Leu Ala	
425 430 435	
TCC TCT TTC AAC TGC ATC TTC CTG TAC ACC GGA GAG CTG TAC CCC	1625
Ser Ser Phe Asn Cys Ile Phe Leu Tyr Thr Gly Glu Leu Tyr Pro	
440 445 450	
ACA GTG ATT CGG CAG ACA GGC CTG GGC ATG GGC AGC ACC ATG GCC	1670
Thr Val Ile Arg Gln Thr Gly Leu Gly Met Gly Ser Thr Met Ala	
455 460 465	
CGG GTG GGC AGC ATT GTG AGC CCG CTG GTG AGC ATG ACT GCA GAG	1715
Arg Val Gly Ser Ile Val Ser Pro Leu Val Ser Met Thr Ala Glu	
470 475 480	
TTC TAC CCC TCC ATG CCT CTC TTC ATC TTC GGC GCT GTC CCT GTG	1760
Phe Tyr Pro Ser Met Pro Leu Phe Ile Phe Gly Ala Val Pro Val	
485 490 495	
GTC GCC AGT GCT GTC ACT GCC CTG CTG CCA GAG ACC TTG GGC CAG	1805
Val Ala Ser Ala Val Thr Ala Leu Leu Pro Glu Thr Leu Gly Gln	
500 505 510	
CCG CTG CCA GAT ACA GTG CAG GAC CTG AAG AGC AGG AGC AGA GGA	1850
Pro Leu Pro Asp Thr Val Gln Asp Leu Lys Ser Arg Ser Arg Gly	
515 520 525	
AAG CAG AAT CAA CAG CAG CAG GAA CAG CAG AAG CAG ATG ATG CCG	1895
Lys Gln Asn Gln Gln Gln Gln Glu Gln Gln Lys Gln Met Met Pro	

530	535	540	
CTC CAG GCC TCA ACA CAA GAG AAG AAT GGA CTT			1928
Leu Gln Ala Ser Thr Gln Glu Lys Asn Gly Leu			
545	550 551		
TGAGAACGGA AGGGCTTCAC ACAGCACTAA AGGGAGTGGG GTTCTACAGG TCCTGCCGTC			1988
TACATGAGGA GGGGGAGTGA GTAGAGGGAC TGGACCATCC AAATGTGGAG GCTGCCATTC			2048
AGAGAAATCC CTCCCCAAAG GTCATGTCAG TAGACCCACT AGGAACAAAA GCTCTGACTA			2108
TGTGCAGCTT CTTAAGCAGA ATGTTCTCGT CACCGGCCAT CTCCTGCTC ATGGTCACTC			2168
CGCCACCTCC AGGACCTTGC AAAGAATCTC AGACAATTAA ATGAATCTCT TCTAAAAAAA			2228
AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA			2288
AAAAAA			2294

[0081]

Sequence ID No: 2

Length: 2171

Type: Nucleic acid

Number of Chain: Doubled-Stranded

Topology: Linear

Kind: cDNA to mRNA

Origin

Organism: Human

Sequence

GAAAGCTGAG CTGCCCTGAC CCCCAAAGTG AGGAGAAGCT GCAAGGGAAA AGGGAGGGAC	60
AGATCAGGGA GACCGGGGAA GAAGGAGGAG CAGCCAAGGA GGCTGCTGTC CCCCCACAGA	120
GCAGCTCGGA CTCAGCTCCC GGAGCAACCC AGCTGCGGAG GCAACGGCAG TGCTGCTCCT	180
CCAGCGAAGG ACAGCAGGCA GGCAGACAGA CAGAGGTCCT GGGACTGGAA GGCCTCAGCC	240

CCCAGCCACT GGGCTGGGCC TGGCCCA	267
ATG GCC TTT AAT GAC CTC CTG CAG CAG GTG GGG GGT GTC GGC CGC	312
Met Ala Phe Asn Asp Leu Leu Gln Gln Val Gly Gly Val Gly Arg	
1 5 10 15	
TTC CAG CAG ATC CAG GTC ACC CTG GTG GTC CTC CCC CTG CTC CTG	357
Phe Gln Gln Ile Gln Val Thr Leu Val Val Leu Pro Leu Leu Leu	
20 25 30	
ATG GCT TCT CAC AAC ACC CTG CAG AAC TTC ACT GCT GCC ATC CCT	402
Met Ala Ser His Asn Thr Leu Gln Asn Phe Thr Ala Ala Ile Pro	
35 40 45	
ACC CAC CAC TGC CGC CCG CCT GCC GAT GCC AAC CTC AGC AAG AAC	447
Thr His His Cys Gly Pro Pro Ala Asp Ala Asn Leu Ser Lys Asn	
50 55 60	
GGG GGG CTG GAG GTC TGG CTG CCC CGG GAC AGG CAG GGG CAG CCT	492
Gly Gly Leu Glu Val Trp Leu Pro Arg Asp Arg Gln Gly Gln Pro	
65 70 75	
GAG TCC TGC CTC CGC TTC ACC TCC CCG CAG TGG GGA CTG CCC TTT	537
Glu Ser Cys Leu Arg Phe Thr Ser Pro Gln Trp Gly Leu Pro Phe	
80 85 90	
CTC AAT GGC ACA GAA GCC AAT GGC ACA GGG GCC ACA GAG CCC TGC	582
Leu Asn Gly Thr Glu Ala Asn Gly Thr Gly Ala Thr Glu Pro Cys	
95 100 105	
ACC GAT GGC TGG ATC TAT GAC AAC AGC ACC TTC CCA TCT ACC ATC	627
Thr Asp Gly Trp Ile Tyr Asp Asn Ser Thr Phe Pro Ser Thr Ile	
110 115 120	

GTG ACT GAG TGG GAC CTT GTG TGC TCT CAC AGG GCC CTA CGC CAG	672
Val Thr Glu Trp Asp Leu Val Cys Ser His Arg Ala Leu Arg Gln	
125 130 135	
CTG GCC CAG TCC TTG TAC ATG GTG GGG GTG CTG CTC GGA GCC ATG	717
Leu Ala Gln Ser Leu Tyr Met Val Gly Val Leu Leu Gly Ala Met	
140 145 150	
GTG TTC GGC TAC CTT GCA GAC AGG CTA GGC CGC CGG AAG GTA CTC	762
Val Phe Gly Tyr Leu Ala Asp Arg Leu Gly Arg Arg Lys Val Leu	
155 160 165	
ATC TTG AAC TAC CTG CAG ACA GCT GTG TCA GGG ACC TGC GCA GCC	807
Ile Leu Asn Tyr Leu Gln Thr Ala Val Ser Gly Thr Cys Ala Arg	
170 175 180	
TTC GCA CCC AAC TTC CCC ATC TAC TGC GCC TTC CGG CTC CTC TCG	852
Phe Ala Pro Asn Phe Pro Ile Tyr Cys Ala Phe Arg Leu Leu Ser	
185 190 195	
GGC ATG GCT CTG GCT GGC ATC TCC CTC AAC TGC ATG ACA CTG AAT	897
Gly Met Ala Leu Ala Gly Ile Ser Leu Asn Cys Met Thr Leu Asn	
200 205 210	
GTG GAG TGG ATG CCC ATT CAC ACA CGG GCC TGC GTG GGC ACC TTG	942
Val Glu Trp Met Pro Ile His Thr Arg Ala Cys Val Gly Thr Leu	
215 220 225	
ATT GGC TAT GTC TAC AGC CTG GGC CAG TTC CTC CTG GCT GGT GTG	987
Ile Gly Tyr Val Tyr Ser Leu Gly Gln Phe Leu Leu Ala Gly Val	
230 235 240	
GCC TAC GCT GTG CCC CAC TGG CGC CAC CTG CAG CTA CTG GTC TCT	1032



Ala Tyr Ala Val Pro His Trp Arg His Leu Gln Leu Leu Val Ser	
245 250 255	
GCG CCT TTT TTT GCC TTC TTC ATC TAC TCC TGG TTC TTC ATT GAG	1077
Ala Pro Phe Phe Ala Phe Phe Ile Tyr Ser Trp Phe Phe Ile Glu	
260 265 270	
TCG GCC CGC TGG CAC TCC TCC TCC GGG AGG CTG GAC CTC ACC CTG	1122
Ser Ala Arg Trp His Ser Ser Ser Gly Arg Leu Asp Leu Thr Leu	
275 280 285	
AGG GCC CTG CAG AGA GTC GCC CGG ATC AAT GGG AAG CGG GAA GAA	1167
Arg Ala Leu Gln Arg Val Ala Arg Ile Asn Gly Lys Arg Glu Glu	
290 295 300	
GGA GCC AAA TTG AGT ATG GAG GTA CTC CGG GCC AGT CTG CAG AAG	1212
Gly Ala Lys Leu Ser Met Glu Val Leu Arg Ala Ser Leu Gln Lys	
305 310 315	
GAG CTG ACC ATG GGC AAA GGC CAG GCA TCG GCC ATG GAG CTG CTG	1257
Glu Leu Thr Met Gly Lys Gly Gln Ala Ser Ala Met Glu Leu Leu	
320 325 330	
CGC TGC CCC ACC CTC CGC CAC CTC TTC CTC TGC CTC TCC ATG CTG	1302
Arg Cys Pro Thr Leu Arg His Leu Phe Leu Cys Leu Ser Met Leu	
335 340 345	
TGG TTT GCC ACT AGC TTT GCA TAC TAT GGG CTG GTC ATG GAC CTG	1347
Trp Phe Ala Thr Ser Phe Ala Tyr Tyr Gly Leu Val Met Asp Leu	
350 355 360	
CAG GGC TTT GGA GTC AGC ATC TAC CTA ATC CAG GTG ATC TTT GGT	1392
Gln Gly Phe Gly Val Ser Ile Tyr Leu Ile Gln Val Ile Phe Gly	

365	370	375	
GCT GTG GAC CTG CCT GCC AAG CTT GTG GGC TTC CTT GTC ATC AAC			1437
Ala Val Asp Leu Pro Ala Lys Leu Val Gly Phe Leu Val Ile Asn			
380	385	390	
TCC CTG GGT CGC CGG CCT GCC CAG ATG GCT GCA CTG CTG CTG GCA			1482
Ser Leu Gly Arg Arg Pro Ala Gln Met Ala Ala Leu Leu Leu Ala			
395	400	405	
GGC ATC TGC ATC CTG CTC AAT GGG GTG ATA CCC CAG GAC CAG TCC			1527
Gly Ile Cys Ile Leu Leu Asn Gly Val Ile Pro Gln Asp Gln Ser			
410	415	420	
ATT GTC CGA ACC TCT CTT GCT GTG CTG GGG AAG GGT TGT CTG GCT			1572
Ile Val Arg Thr Ser Leu Ala Val Leu Gly Lys Gly Cys Leu Ala			
425	430	435	
GCC TCC TTC AAC TGC ATC TTC CTG TAT ACT GGG GAA CTG TAT CCC			1617
Ala Ser Phe Asn Cys Ile Phe Leu Tyr Thr Gly Glu Leu Tyr Pro			
440	445	450	
ACA ATG ATC CGG CAG ACA GGC ATG GGA ATG GGC AGC ACC ATG GCC			1662
Thr Met Ile Arg Gln Thr Gly Met Gly Met Gly Ser Thr Met Ala			
455	460	465	
CGA GTG GGC AGC ATC GTG AGC CCA CTG GTG AGC ATG ACT GCC GAG			1707
Arg Val Gly Ser Ile Val Ser Pro Leu Val Ser Met Thr Ala Glu			
470	475	480	
CTC TAC CCC TCC ATG CCT CTC TTC ATC TAC GGT GCT GTT CCT GTG			1752
Leu Tyr Pro Ser Met Pro Leu Phe Ile Tyr Gly Ala Val Pro Val			
485	490	495	

GCC GCC AGC GCT GTC ACT GTC CTC CTG CCA GAG ACC CTG GGC CAG	1797
Ala Ala Ser Ala Val Thr Val Leu Leu Pro Glu Thr Leu Gly Gln	
500 505 510	
CCA CTG CCA GAC ACG GTG CAG GAC CTG GAG AGC AGG TGG GCC CCC	1842
Pro Leu Pro Asp Thr Val Gln Asp Leu Glu Ser Arg Trp Ala Pro	
515 520 525	
ACT CAG AAA GAA GCA GGG ATA TAT CCC AGG AAA GGG AAA CAG ACG	1887
Thr Gln Lys Glu Ala Gly Ile Tyr Pro Arg Lys Gly Lys Gln Thr	
530 535 540	
CGA CAG CAA CAA GAG CAC CAG AAG TAT ATG GTC CCA CTG CAG GCC	1932
Arg Gln Gln Gln Glu His Gln Lys Tyr Met Val Pro Leu Gln Ala	
545 550 555	
TCA GCA CAA GAG AAG AAT GGA CTC	1956
Ser Ala Gln Glu Lys Asn Gly Leu	
560 563	
TGAGGACTGA GAAGGGGCCT TACAGAACCC TAAAGGGAGG GAAGGTCCTA CAGGTCTCCG	2016
GCCACCCACA CAAGGAGGAG GAAGAGGAAA TGGTGACCCA AGTGTGGGGG TTGTGGTTCA	2076
GGAAAGCATC TTCCAGGGG TCCACCTCCC TTTATAAACC CCACCAGAAC CACATCATTA	2136
AAAGGTTTGA CTGCGAAAAA AAAAAAAAAA AAAAA	2171

[Brief Description of the Drawings]

[Fig. 1] This is a drawing which shows the result of an uptake experiment of glutaric acid by oocytes into which cRNA of sodium-dependent dicarboxylate transporter (NaDC-1) gene of rat was injected.

[Fig. 2] This is a drawing which shows the result of an uptake experiment of PAH by oocytes into which mRNA derived from renal tissues of rat and/or cRNA of NaDC-1 gene of rat were/was injected.

[Fig. 3] This is a drawing which shows hydrophobic plots of rat organic anion transporter OAT1.

[Fig. 4] This is a photographic picture which shows the result where expression of OAT1 gene mRNA in various organ tissues of rat was analyzed by a northern blotting.

[Fig. 5] This is a drawing which shows the result where influence of preincubation with glutaric acid was investigated in a PAH uptake experiment by oocytes into which rat OAT1 gene cRNA was injected.


[Fig. 6] This is a drawing which shows the result of investigation of influence of salt added in a PAH uptake experiment by oocytes into which rat OAT1 gene cRNA was injected.

[Fig. 7] This is a drawing which shows the result of investigation of influence of concentration of the substrate PAH in a PAH uptake experiment by oocytes into which rat OAT1 gene cRNA was injected.


[Fig. 8] This is a drawing which shows the result of investigation of influence of addition of various drugs to the system in a PAH uptake experiment by oocytes into which rat OAT1 gene cRNA was injected.

[Fig. 9] This is a drawing which shows the result of an uptake experiment by oocytes into which cRNA of rat OAT1 gene was injected when various drugs were used as substrates.


[Figure 1]



[Figure 2]




[Figure 3]







[Figure 4]




【Figure 5】




[Figure 6]




[Figure 7]



[Figure 8]





[Figure 9]

[Designation of Document] Abstract

[Abstract]

[Problem] Novel organic anion transporter and gene coding thereof are provided.

[Means for Resolution] Protein having a capability of transporting an organic anion and comprising an amino acid sequence represented by SEQ ID No. 1 or 2 or an amino acid sequence represented by SEQ ID No. 1 or 2 where one or several amino acid(s) is/are deleted therefrom, substituted therefor or added thereto. Gene coding the above-mentioned protein.

[Selected Drawing] (none)